

**IMPROVED MUSCLE-DIRECTED GENE TRANSFER BY USE OF
RECOMBINANT AAV-1 AND AAV-6 VIRIONS**

5 **CROSS REFERENCE TO RELATED APPLICATIONS**

This application claims priority under 37 C.F.R. § 119(e) to Provisional Application Ser. No. 60/266,778 filed on February 6, 2001, herein incorporated by reference in its entirety.

10

FIELD OF THE INVENTION

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
218

in order to maintain the necessary levels of the protein in the bloodstream. Even then, the concentration of protein is not maintained at physiological levels—the level of the protein is usually abnormally high immediately following the injection, and far below optimal levels prior to the injection. Additionally, injected delivery of recombinant protein often cannot deliver the protein to the target cells, tissues, or organs in the body. And, if the protein successfully reaches its target, it may be diluted to a non-therapeutic level. Furthermore, the method is inconvenient and often restricts the patient's lifestyle.

These shortcomings have fueled the desire to develop gene therapy methods for delivering sustained levels of specific proteins into patients. These methods are designed to allow clinicians to introduce deoxyribonucleic acid (DNA) coding for a nucleic acid, such as a therapeutic gene, directly into a patient (*in vivo* gene therapy) or into cells isolated from a patient or a donor (*ex vivo* gene therapy). The introduced nucleic acid then directs the patient's own cells or grafted cells to produce the desired protein product. Gene delivery, therefore, obviates the need for frequent injections. Gene therapy may also allow clinicians to select specific organs or cellular targets (e.g., muscle, blood cells, brain cells, etc.) for therapy.

DNA may be introduced into a patient's cells in several ways. There are transfection methods, including chemical methods such as calcium phosphate precipitation and liposome-mediated transfection, and physical methods such as electroporation. In general, transfection methods are not suitable for *in vivo* gene delivery. There are also methods that use recombinant viruses. Current viral-mediated gene delivery vectors include those based on retrovirus, adenovirus, herpes virus, pox virus, and adeno-associated virus (AAV). Like the retroviruses, and unlike adenovirus, AAV has the ability to integrate its genome into a host cell chromosome.

Adeno-Associated Virus-Mediated Gene Therapy

AAV is a parvovirus belonging to the genus Dependovirus, and has several attractive features not found in other viruses. For example, AAV can infect a wide range of host cells, including non-dividing cells. AAV can also infect cells from different species. Importantly, AAV has not been associated with any human or animal

disease, and does not appear to alter the physiological properties of the host cell upon integration. Furthermore, AAV is stable at a wide range of physical and chemical conditions, which lends itself to production, storage, and transportation requirements.

The AAV genome, a linear, single-stranded DNA molecule containing approximately 4700 nucleotides (the AAV-2 genome consists of 4681 nucleotides), generally comprises an internal non-repeating segment flanked on each end by inverted terminal repeats (ITRs). The ITRs are approximately 145 nucleotides in length (AAV-1 has ITRs of 143 nucleotides) and have multiple functions, including serving as origins of replication, and as packaging signals for the viral genome.

The internal non-repeated portion of the genome includes two large open reading frames (ORFs), known as the AAV replication (*rep*) and capsid (*cap*) regions. These ORFs encode replication and capsid gene products, respectively: replication and capsid gene products (i.e., proteins) allow for the replication, assembly, and packaging of a complete AAV virion. More specifically, a family of at least four viral proteins are expressed from the AAV *rep* region: Rep 78, Rep 68, Rep 52, and Rep 40, all of which are named for their apparent molecular weights. The AAV *cap* region encodes at least three proteins: VP1, VP2, and VP3.

In nature, AAV is a helper virus-dependent virus, i.e., it requires co-infection with a helper virus (e.g., adenovirus, herpesvirus, or vaccinia virus) in order to form functionally complete AAV virions. In the absence of co-infection with a helper virus, AAV establishes a latent state in which the viral genome inserts into a host cell chromosome or exists in an episomal form, but infectious virions are not produced. Subsequent infection by a helper virus “rescues” the integrated genome, allowing it to be replicated and packaged into viral capsids, thereby reconstituting the infectious virion. While AAV can infect cells from different species, the helper virus must be of the same species as the host cell. Thus, for example, human AAV will replicate in canine cells that have been co-infected with a canine adenovirus.

To construct infectious recombinant AAV (rAAV) containing a nucleic acid, a suitable host cell line is transfected with an AAV vector containing a nucleic acid. AAV helper functions and accessory functions are then expressed in the host cell. Once these factors come together, the HNA is replicated and packaged as though it were a

wild-type (wt) AAV genome, forming a recombinant virion. When a patient's cells are infected with the resulting rAAV, the HNA enters and is expressed in the patient's cells. Because the patient's cells lack the *rep* and *cap* genes, as well as the adenovirus accessory function genes, the rAAV are replication defective; that is, they cannot further replicate and package their genomes. Similarly, without a source of *rep* and *cap* genes, wtAAV cannot be formed in the patient's cells.

There are six known AAV serotypes, AAV-1 through AAV-6. Of those six serotypes, AAV-2 is the best characterized, having been used to successfully deliver transgenes to several cell lines, tissue types, and organs in a variety of in vitro and in vivo assays. The six serotypes of AAV can be distinguished from one another by the use of monoclonal antibodies or by employing nucleotide sequence analysis; AAV-1, AAV-2, AAV-3, and AAV-6 are 82% identical at the nucleotide level, while AAV-4 is 75 to 78% identical to the other serotypes (Russell et al. (1998) *J Virol* 72:309-319). Significant nucleotide sequence variation is noted for regions of the AAV genome that code for capsid proteins; such variable regions may be responsible for differences in serological reactivity to the capsid proteins of the various AAV serotypes.

It is known that readministration of a single AAV serotype can lead to a significant reduction in transduction efficiency. Moskalenko et al. (*J Virol* (2000) 74:176101766), for example, showed that mice with pre-existing anti-AAV-2 antibodies, when administered Factor IX in a recombinant AAV-2 virion, failed to express the Factor IX transgene, suggesting that the anti-AAV-2 antibodies blocked transduction of the rAAV-2 virion. Halbert et al. (*J Virol* (1998) 72:9795-9805) reported similar results. Others have demonstrated successful readministration of rAAV-2 virions into experimental animals, but only after immune suppression is achieved (e.g., Halbert et al., supra).

Thus, using rAAV-2 for human gene therapy is potentially problematic because anti-AAV-2 antibodies are prevalent in human populations; in fact, one study estimated that at least 80% of the general population has been infected with AAV-2 (Berns and Linden (1995) *Bioessays* 17:237-245). The identification of AAV serotypes that are not serologically cross-reactive with AAV-2 would be a significant advancement in the art. Such AAV serotypes are described herein.

SUMMARY OF THE INVENTION

The present invention provides AAV serotypes that have the ability to efficiently transduce cell and tissue types that AAV-2 transduces poorly and/or will not be inhibited by anti-AAV-2 antibodies. In accordance with the present invention, methods and AAV vectors for use therein are provided for the efficient delivery of a heterologous nucleic acid molecule(s) (HNA) to cells or tissue of a mammal, using recombinant AAV virions. Preferably, the cells or tissue are muscle cells or muscle tissue.

More specifically, the present invention provides for the use of AAV-1 and AAV-6 serotypes (i.e., AAV virions containing AAV-1 and/or AAV-6 capsid proteins) to deliver an HNA encoding anti-sense RNA, ribozymes, or genes that express proteins, wherein expression of said anti-sense RNA, ribozymes, or genes provides for a therapeutic effect in a mammalian subject. In one embodiment, the rAAV virions containing an HNA are injected directly into a muscle. In another embodiment, the rAAV virions containing an HNA are administered into the vasculature via injection into veins, arteries, or other vascular conduits, or by using techniques such as isolated limb perfusion.

In a preferred embodiment of the invention, AAV-1-derived and AAV-6-derived virions are provided that contain a gene encoding a blood coagulation protein which, when expressed at a sufficient concentration, provides for a therapeutic effect, such effect being an improvement in the blood-clotting efficiency of a mammal suffering from a blood clotting disorder. The blood clotting disorder can be any disorder adversely affecting the organism's ability to coagulate the blood. Preferably, the blood clotting disorder is hemophilia.

In one embodiment, the gene encoding a blood coagulation protein is a Factor VIII gene. Preferably, the Factor VIII gene is the human Factor VIII gene or a derivation thereof. In another embodiment, the gene encoding a blood coagulation protein is a Factor IX gene. Preferably, the Factor IX gene is the human Factor IX (hF.IX) gene.

These and other embodiments of the instant invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

BRIEF DESCRIPTION OF THE DRAWINGS

5

FIG. 1 represents circulating plasma hF.IX in nanograms per milliliter (ng/mL) as measured in RAG-1 mice following intramuscular (IM) injection of 2×10^{11} viral vector genomes/kg (n=4).

10

FIG. 1 represents circulating plasma hF.IX in nanograms per milliliter (ng/mL) as measured in RAG-1 mice following intramuscular (IM) injection of 2×10^{11} viral vector genomes/kg (n=4).

DETAILED DESCRIPTION OF THE INVENTION

The present invention embraces the use of a recombinant adeno-associated virus (rAAV) virion to deliver a “heterologous nucleic acid” (an “HNA”) to a mammalian subject. A “recombinant AAV virion” or “rAAV virion” is an infectious virus composed of an AAV protein shell (i.e., a capsid) encapsulating a “recombinant AAV (rAAV) vector,” the rAAV vector comprising the HNA and one or more AAV inverted terminal repeats (ITRs). AAV vectors can be constructed using recombinant techniques that are known in the art and include one or more HNAs flanked by functional ITRs. The ITRs of the rAAV vector need not be the wild-type nucleotide sequences, and may be altered, e.g., by the insertion, deletion, or substitution of nucleotides, so long as the sequences provide for proper function, i.e., rescue, replication, and packaging of the AAV genome.

Recombinant AAV virions may be produced using a variety of techniques known in the art, including the triple transfection method (described in detail in U.S. Patent No. 6,001,650, the entirety of which is incorporated by reference). This system involves the use of three vectors for rAAV virion production, including an AAV helper function vector, an accessory function vector, and a rAAV vector that contains the HNA. One of skill in the art will appreciate, however, that the nucleic acid sequences encoded by these vectors can be provided on two or more vectors in various combinations. As used herein, the term “vector” includes any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, artificial chromosome, virus, virion, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

The AAV helper function vector encodes the “AAV helper function” sequences (i.e., rep and cap), which function *in trans* for productive AAV replication and encapsidation. Preferably, the AAV helper function vector supports efficient AAV vector production without generating any detectable wild-type AAV virions (i.e., AAV virions containing functional rep and cap genes). An example of such a vector, pHLP19 is described in U.S. Patent No. 6,001,650. Another AAV helper function vector is the

pRep6cap6 vector, described in U.S. Patent No. 6,156,303, the entirety of which is herein incorporated by reference.

The accessory function vector encodes nucleotide sequences for non-AAV derived viral and/or cellular functions upon which AAV is dependent for replication (i.e., “accessory functions”). The accessory functions include those functions required for AAV replication, including, without limitation, those moieties involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of cap expression products, and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses such as adenovirus, herpesvirus (other than herpes simplex virus type-1), and vaccinia virus. In a preferred embodiment, the accessory function plasmid pladen05 is used (details regarding pLaden05 are described in U.S. Patent No. 6,004,797, the entirety of which is hereby incorporated by reference). This plasmid provides a complete set of adenovirus accessory functions for AAV vector production, but lacks the components necessary to form replication-competent adenovirus.

The instant invention broadly contemplates the use of two specific AAV serotypes: AAV-1 and the AAV-6. The term “serotype” is used herein to describe the genotype of a virus or unicellular organism that has been defined by means of antisera binding to antigenic determinants located on the surface of the virus or unicellular organism. In the case of AAV, the antigenic determinants located on the AAV virion’s surface are the capsid proteins, so it is the capsid protein that distinguishes the AAV serotype. Capsid proteins are the product of AAV cap gene expression, the specific sequence of the cap gene being unique to the particular AAV serotype.

Recombinant AAV-6 virions are produced, in one embodiment of the invention, with an AAV helper function vector containing the rep and cap genes from the AAV-6 genome (this vector is known as pRepCap6 – see U.S. Patent No. 6,156,303, supra, for a thorough description of the pRepCap6 vector). The AAV-6 genome has been published and is available under GenBank Accession No. 9629894. One of skill in the art will also appreciate that other rep genes, e.g. rep2, can be used in combination with the cap6 gene to produce AAV “hybrid” helper function vectors, which are then capable of supporting the production of rAAV-6 virions. The term “hybrid” as used herein is an

AAV helper function vector with a rep gene from one serotype (other than AAV-6) in combination with the cap gene from the AAV-6 genome (e.g., pRep2Cap6, pRep3Cap6, etc.). The rep and cap genes can be wild-type in their sequences, or be altered by partial deletion, mutation, rearrangement, addition, gene or gene segment shuffling, etc., the primary consideration as contemplated herein being the retention of rep and cap wild-type function. See U.S. Patent No. 6,156,303, *supra*, (and Example 1 below) for methods describing the generation of rAAV-6 virions.

Similar methods can also be employed to construct AAV helper function vectors containing rep and cap1 genes. Incorporating the rep gene from the AAV-1 genome with the cap gene from the AAV-1 genome yields the AAV helper function vector pRepCap1. Incorporating a rep gene from an AAV serotype other than AAV-1 yields a hybrid AAV helper function vector still capable of supporting AAV-1 virion production since the AAV helper function vector contains the cap1 gene. Either pRepCap1 or a hybrid AAV helper function vector such as pRep2Cap1 can support rAAV-1 virion production. The AAV-1 genome has been published under the Patent Cooperation Treaty (international publication WO 0028061) and is available under GenBank Accession No. 9632547.

We have shown that the cap1 and cap6 proteins recognize the cap binding site(s) on the AAV-2 ITRs. This is thought to be because the cap1 and cap6 proteins recognize the AAV-2 ITR secondary structure, and not specific AAV-2 ITR DNA sequences. It is believed that the ITRs from the various AAV serotypes assume similar secondary structure so one of skill in the art would appreciate that AAV-1, AAV-3, AAV-4, AAV-5, or AAV-6 ITRs could be used with pRep6Cap6 or a hybrid AAV-6 helper function vector to generate AAV-6 serotype virions (or pRep1Cap1 or a hybrid AAV-1 helper function vector to generate AAV-1 serotype virions). For example, using the methods of the instant invention, pRep2Cap6 could be used in conjunction with AAV-3 ITRs to produce AAV-6 virions.

The HNA, that is, the "heterologous nucleic acid," comprises nucleic acid sequences joined together that are otherwise not found together in nature, this concept defining the term "heterologous." To illustrate the point, an example of an HNA is a gene flanked by nucleotide sequences not found in association with that gene in nature.

Another example of an HNA is a gene that itself is not found in nature (e.g., synthetic sequences having codons different from the native gene). Allelic variation or naturally occurring mutational events do not give rise to HNAs, as used herein. An HNA can comprise an anti-sense RNA molecule, a ribozyme, or a gene encoding a polypeptide.

5 The HNA is operably linked to a heterologous promoter (constitutive, cell-specific, or inducible) such that the HNA is capable of being expressed in the patient's target cells under appropriate or desirable conditions. Numerous examples of constitutive, cell-specific, and inducible promoters are known in the art, and one of skill could readily select a promoter for a specific intended use, e.g., the selection of the
10 muscle-specific skeletal α -actin promoter or the muscle-specific creatine kinase promoter/enhancer for muscle cell-specific expression, the selection of the constitutive CMV promoter for strong levels of continuous or near-continuous expression, or the selection of the inducible ecdysone promoter for induced expression. Induced expression allows the skilled artisan to control the amount of protein that is synthesized.
15 In this manner, it is possible to vary the concentration of therapeutic product. Other examples of well known inducible promoters are: steroid promoters (e.g., estrogen and androgen promoters) and metallothionein promoters.

 The invention includes rAAV-1 or rAAV-6 virions comprising HNAs coding for one or more anti-sense RNA molecules, the rAAV virions preferably administered to
20 one or more muscle cells or tissue of a mammal. Antisense RNA molecules suitable for use with the present invention in cancer anti-sense therapy or treatment of viral diseases have been described in the art. See, e.g., Han et al., (1991) *Proc. Natl. Acad. Sci. USA* 88:4313-4317; Uhlmann et al., (1990) *Chem. Rev.* 90:543-584; Helene et al., (1990) *Biochim. Biophys. Acta.* 1049:99-125; Agarawal et al., (1988) *Proc. Natl. Acad. Sci. USA* 85:7079-7083; and Heikkila et al., (1987) *Nature* 328:445-449. The invention also
25 encompasses the delivery of ribozymes using the methods disclosed herein. For a discussion of suitable ribozymes, see, e.g., Cech et al., (1992) *J. Biol. Chem.* 267:17479-17482 and U.S. Pat. No. 5,225,347.

 The invention preferably encompasses rAAV-1 or rAAV-6 virions comprising
30 HNAs coding for one or more polypeptides, the rAAV virions preferably administered to one or more muscle cells or tissue of a mammal. Thus, the invention embraces the

delivery of HNAs encoding one or more peptides, polypeptides, or proteins, which are useful for the treatment of disease states in a mammalian subject. Such DNA and associated disease states include, but are not limited to: DNA encoding glucose-6-phosphatase, associated with glycogen storage deficiency type 1A; DNA encoding phosphoenolpyruvate-carboxykinase, associated with Pepck deficiency; DNA encoding galactose-1 phosphate uridyl transferase, associated with galactosemia; DNA encoding phenylalanine hydroxylase, associated with phenylketonuria; DNA encoding branched chain alpha-ketoacid dehydrogenase, associated with Maple syrup urine disease; DNA encoding fumarylacetoacetate hydrolase, associated with tyrosinemia type 1; DNA encoding methylmalonyl-CoA mutase, associated with methylmalonic acidemia; DNA encoding medium chain acyl CoA dehydrogenase, associated with medium chain acetyl CoA deficiency; DNA encoding ornithine transcarbamylase, associated with ornithine transcarbamylase deficiency; DNA encoding argininosuccinic acid synthetase, associated with citrullinemia; DNA encoding low density lipoprotein receptor protein, associated with familial hypercholesterolemia; DNA encoding UDP-glucouronosyltransferase, associated with Crigler-Najjar disease; DNA encoding adenosine deaminase, associated with severe combined immunodeficiency disease; DNA encoding hypoxanthine guanine phosphoribosyl transferase, associated with Gout and Lesch-Nyan syndrome; DNA encoding biotinidase, associated with biotinidase deficiency; DNA encoding beta-glucocerebrosidase, associated with Gaucher disease; DNA encoding beta-glucuronidase, associated with Sly syndrome; DNA encoding peroxisome membrane protein 70 kDa, associated with Zellweger syndrome; DNA encoding porphobilinogen deaminase, associated with acute intermittent porphyria; DNA encoding alpha-1 antitrypsin for treatment of alpha-1 antitrypsin deficiency (emphysema); DNA encoding erythropoietin for treatment of anemia due to thalassemia or to renal failure; DNA encoding vascular endothelial growth factor, DNA encoding angiopoietin-1, and DNA encoding fibroblast growth factor for the treatment of ischemic diseases; DNA encoding thrombomodulin and tissue factor pathway inhibitor for the treatment of occluded blood vessels as seen in, for example, atherosclerosis, thrombosis, or embolisms; DNA encoding aromatic amino acid decarboxylase (AADC), and DNA encoding tyrosine hydroxylase (TH) for the treatment of Parkinson's disease;

DNA encoding the beta adrenergic receptor, DNA encoding anti-sense to, or DNA encoding a mutant form of, phospholamban, DNA encoding the sarco(endo)plasmic reticulum adenosine triphosphatase-2 (SERCA2), and DNA encoding the cardiac adenylyl cyclase for the treatment of congestive heart failure; DNA encoding a tumor suppressor gene such as p53 for the treatment of various cancers; DNA encoding a cytokine such as one of the various interleukins for the treatment of inflammatory and immune disorders and cancers; DNA encoding dystrophin or minidystrophin and DNA encoding utrophin or miniutrophin for the treatment of muscular dystrophies; and, DNA encoding insulin for the treatment of diabetes.

The invention also includes rAAV-1 and rAAV-6 virions comprising a gene or genes coding for blood coagulation proteins, which proteins may be delivered, using the methods of the present invention, to the cells of a mammal having hemophilia for the treatment of hemophilia. Thus, the invention includes: delivery of the Factor IX gene to a mammal for treatment of hemophilia B, delivery of the Factor VIII gene to a mammal for treatment of hemophilia A, delivery of the Factor VII gene for treatment of Factor VII deficiency, delivery of the Factor X gene for treatment of Factor X deficiency, delivery of the Factor XI gene for treatment of Factor XI deficiency, delivery of the Factor XIII gene for treatment of Factor XIII deficiency, and, delivery of the Protein C gene for treatment of Protein C deficiency. Delivery of each of the above-recited genes to the cells of a mammal is accomplished by first generating a rAAV virion comprising the gene and then administering the rAAV virion to the mammal. Thus, the invention includes rAAV virions comprising genes encoding any one of Factor IX, Factor VIII, Factor X, Factor VII, Factor XI, Factor XIII or Protein C.

Delivery of rAAV-1 or rAAV-6 virions containing one or more HNAs to a mammalian subject may be by intramuscular injection or by administration into the bloodstream of the mammalian subject. Administration into the bloodstream may be by injection into a vein, an artery, or any other vascular conduit such as a venule, an arteriole, or capillary. Additionally, a skilled artisan can administer rAAV-1 or rAAV-6 virions into the bloodstream by way of isolated limb perfusion, a technique well known in the surgical arts, the method essentially enabling the artisan to isolate a limb from the systemic circulation prior to administration of the rAAV virions. A variant of the

isolated limb perfusion technique, described in U.S. Patent No. 6,177,403 and herein incorporated by reference, can also be employed by the skilled artisan to administer rAAV-1 or rAAV-6 virions into the vasculature of an isolated limb to potentially enhance transduction into muscle cells or tissue.

5 The dose of rAAV virions required to achieve a particular “therapeutic effect,” e.g., the units of dose in vector genomes/per kilogram of body weight (vg/kg), will vary based on several factors including, but not limited to: the route of rAAV virion administration, the level of gene (or anti-sense RNA or ribozyme) expression required to achieve a therapeutic effect, the specific disease or disorder being treated, a host
10 immune response to the rAAV virion, a host immune response to the gene (or anti-sense RNA or ribozyme) expression product, and the stability of the gene (or anti-sense RNA or ribozyme) product. One of skill in the art can readily determine a rAAV virion dose range to treat a patient having a particular disease or disorder based on the aforementioned factors, as well as other factors that are well known in the art.

15 Generally speaking, by “therapeutic effect” is meant a level of expression of one or more HNAs sufficient to alter a component of a disease (or disorder) toward a desired outcome or clinical endpoint, such that a patient’s disease or disorder shows clinical improvement, often reflected by the amelioration of a clinical sign or symptom relating to the disease or disorder. Using hemophilia as a specific disease example, a
20 “therapeutic effect” for hemophilia is defined herein as an increase in the blood-clotting efficiency of a mammal afflicted with hemophilia, efficiency being determined, for example, by well known endpoints or techniques such as employing assays to measure whole blood clotting time or activated prothromboplastin time. Reductions in either whole blood clotting time or activated prothromboplastin time are indications of an
25 increase in blood-clotting efficiency. In severe cases of hemophilia, hemophiliacs having less than 1% of normal levels of Factor VIII or Factor IX have a whole blood clotting time of greater than 60 minutes as compared to approximately 10 minutes for non-hemophiliacs. Expression of 1% or greater of Factor VIII or Factor IX has been shown to reduce whole blood clotting time in animal models of hemophilia, so
30 achieving a circulating Factor VIII or Factor IX plasma concentration of greater than

1% will likely achieve the desired therapeutic effect of an increase in blood-clotting efficiency.

Rather than focusing exclusively on treating a disease, it is often desirable to deliver an HNA to a host cell in order to elucidate its physiological or biochemical function(s). The HNA can be either an endogenous gene or heterologous. Using either an ex vivo or in vivo approach, the skilled artisan can administer rAAV-1 and/or rAAV-6 virions containing one or more HNAs of unknown function to an experimental animal, express the HNA(s), and observe any subsequent functional changes. Such changes can include: protein-protein interactions, alterations in biochemical pathways, alterations in the physiological functioning of cells, tissues, organs, or organ systems, and/or the stimulation or silencing of gene expression.

Alternatively, the skilled artisan can over-express a gene of known function and examine its effects. Such genes can be either endogenous to the experimental animal or heterologous in nature (i.e., a transgene).

By using the methods of the present invention, the skilled artisan can also abolish or significantly reduce gene expression, thereby employing another means of determining gene function. One method of accomplishing this is by way of administering antisense RNA-containing rAAV virions to an experimental animal, expressing the antisense RNA molecule so that the targeted endogenous gene is “knocked out,” and then observing any subsequent physiological or biochemical changes.

The methods of the present invention are compatible with other well-known technologies such as transgenic mice and knockout mice and can be used to complement these technologies. One skilled in the art can readily determine combinations of known technologies with the methods of the present invention to obtain useful information on gene function.

Once delivered, in many instances it is not enough to simply express the HNA; instead, it is often desirable to vary the levels of HNA expression. Varying HNA expression levels, which varies the dose of the HNA expression product, is frequently useful in acquiring and/or refining functional information on the HNA. This can be accomplished, for example by incorporating a heterologous inducible promoter into the

5 rAAV virion containing the HNA so that the HNA will be expressed only when the promoter is induced. Some inducible promoters can also provide the capability for refining HNA expression levels; that is, varying the concentration of inducer will fine-tune the concentration of HNA expression product. This is sometimes more useful than having an "on-off" system (i.e., any amount of inducer will provide the same level of HNA expression product, an "all or none" response). Numerous examples of inducible promoters are known in the art including the ecdysone promoter, steroid promoters (e.g., estrogen and androgen promoters) and metallothionein promoters.

10 The methods of the present invention can be used to facilitate pharmaco- or toxico-kinetic studies. For example, because AAV is known to transduce hepatocytes with high efficiency, human metabolic enzymes (e.g., various oxidases and reductases such as the cytochrome p450 isozymes, various epoxide hydrolases, various dehydrogenases such as alcohol and aldehyde dehydrogenases, various peptidases, etc. – metabolic enzymes that are expressed and function in hepatocytes) can be delivered to the liver of mice by way of rAAV-1 and or rAAV-6 virions, expressed, and then various drugs and/or toxicants can be administered to the transduced mice in order to screen for any metabolites of interest.

15 The presented methods resulted in an unexpected transduction efficiency of rAAV-6 in the skeletal muscle of mice, with transduction efficiency measured by circulating plasma levels of hF.IX (the hF.IX gene delivered to the skeletal muscle of mice by rAAV-6 virions). As shown in FIG. 1, after three weeks post-injection of rAAV-6-hF.IX, serum levels of hF.IX were approximately 32-fold greater than serum levels of hF.IX in mice injected with rAAV2-hF.IX virions. After seven weeks following injection, hF.IX delivered by rAAV-6 remained at higher concentrations than rAAV-2-delivered hF.IX. For rAAV-1-hF.IX, circulating hF.IX levels were approximately 18-fold higher than circulating F.IX levels obtained from rAAV-2-hF.IX mice. After eleven weeks post-injection, the difference between rAAV-1-hF.IX and rAAV-2-hF.IX increased to 50-fold.

20 The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

EXAMPLE 1
RECOMBINANT AAV FACTOR IX VIRION PREPARATION

5 Recombinant AAV virions containing the human Factor IX (hF.IX) gene – the complete cDNA sequence for hF.IX available under GenBank Accession No. 182612 – were prepared using a triple-transfection procedure described in U.S. Patent No. 6,001,650, supra.

Vector Construction

AAV pRepCap6 Helper Function Vector

10 The pRepCap6 AAV helper function vector was constructed using standard molecular biological techniques. Using an infectious AAV-6 virion (the published wild-type sequence available under GenBank Accession No. 9629894), two Bgl II restriction sites were engineered into the AAV-6 genome, one just upstream of the p5
15 promoter and one downstream of the polyadenylation site, creating the pAAV6Bgl plasmid. The Bgl II fragment containing the rep6 and cap6 genes was excised from pAAV6Bgl and inserted into a pBLUE-SCRIPT (Stratagene, La Jolla, CA) backbone to create the AAV-6 helper vector pRepCap6.

pLadeno5 Accessory Function Vector

20 The accessory function vector pLadeno5 was constructed as follows: DNA fragments encoding the E2a, E4, and VA RNA regions isolated from purified adenovirus serotype-2 DNA (obtained from Gibco/BRL) were ligated into a plasmid called pAmpscript. The pAmpscript plasmid was assembled as follows: oligonucleotide-directed mutagenesis was used to eliminate a 623-bp region including
25 the polylinker and alpha complementation expression cassette from pBSII s/k+ (obtained from Stratagene), and replaced with an EcoRV site. The sequence of the mutagenic oligo used on the oligonucleotide-directed mutagenesis was 5'-CCGCTACAGGGCGCGATATCAGCTCACTCAA-3'. A polylinker (containing the following restriction sites: Bam HI; KpnI; SrfI; XbaI; ClaI; Bst1107I; SalI; PmeI; and
30 NdeI) was synthesized and inserted into the EcoRV site created above such that the BamHI side of the linker was proximal to the fl origin in the modified plasmid to

provide the pAmpscript plasmid. The sequence of the polylinker was 5'-GGATCCGGTACCGCCCGGGCTCTAGAATCGATGTATACGTCGACGTTTAAACCATATG-3'.

DNA fragments comprising the adenovirus serotype-2 E2a and VA RNA sequences were cloned directly into pAmpscript. In particular, a 5962-bp SrfI-KpnI(partial) fragment containing the E2a region was cloned between the SrfI and KpnI sites of pAmpscript. The 5962-bp fragment comprises base pairs 21,606-27,568 of the adenovirus serotype-2 genome. The complete sequence of the adenovirus serotype-2 genome is accessible under GenBank No. 9626158.

The DNA comprising the adenovirus serotype-2 E4 sequences had to be modified before it could be inserted into the pAmpscript polylinker. Specifically, PCR mutagenesis was used to replace the E4 proximal, adenoviral terminal repeat with a SrfI site. The location of this SrfI site is equivalent to base pairs 35,836-35,844 of the adenovirus serotype-2 genome. The sequences of the oligonucleotides used in the mutagenesis were: 5'-AGAGGCCCGGGCGTTTTAGGGCGGAGTAACTTGC-3' and 5'-ACATACCCGCAGGCGTAGAGAC-3'. A 3,192 bp E4 fragment, produced by cleaving the above-described modified E4 gene with SrfI and SpeI, was ligated between the SrfI and XbaI sites of pAmpscript which already contained the E2a and VA RNA sequences to result in the pLaden5 plasmid. The 3,192-bp fragment is equivalent to base pairs 32,644-35,836 of the adenovirus serotype-2 genome.

rAAV-2 hF.IX vector

The rAAV-2 hF.IX vector is an 11,442-bp plasmid containing the cytomegalovirus (CMV) immediate early promoter, exon 1 of hF.IX, a 1.4-kb fragment of hF.IX intron 1, exons 2-8 of h.FIX, 227 bp of h.FIX 3' UTR, and the SV40 late polyadenylation sequence between the two AAV-2 inverted terminal repeats (U.S. Patent No. 6,093,392, herein incorporated by reference). The 1.4-kb fragment of hF.IX intron 1 consists of the 5' end of intron 1 up to nucleotide 1098 and the sequence from nucleotide 5882 extending to the junction with exon 2. The CMV immediate early promoter and the SV40 late polyadenylation signal sequences can be obtained from the published sequence of pCMV-Script®, which is available from the Stratagene catalog, Stratagene, La Jolla, CA, and from their website, www.stratagene.com.

Triple Transfection Procedure

Specifically, for rAAV-6 virion production, the AAV helper function rep⁶cap⁶ vector (described in U.S. Patent No. 6,156,303, supra), the accessory function vector pLadeno5 (described in U.S. Patent No. 6,004,797, supra), and the rAAV2-hF.IX vector (U.S. Patent No. 6,093,392, supra) were used. Briefly, human embryonic kidney cells type 293 (293 cells – available from the American Type Culture Collection, catalog number CRL-1573) were seeded in 10 cm tissue culture-treated sterile dishes at a density of 3×10^6 cells per dish in 10 mL of cell culture medium consisting of Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum and incubated in a humidified environment at 37° C in 5% CO₂. After overnight incubation, 293 cells were approximately eighty-percent confluent. The 293 cells were then transfected with DNA by the calcium phosphate precipitate method, a transfection method well known in the art. Briefly, 10 µg of each vector (pRepCap6, pLadeno5, and rAAV2-hF.IX) were added to a 3-mL sterile, polystyrene snap cap tube using sterile pipette tips. 1.0 mL of 300 mM CaCl₂ (JRH grade) was added to each tube and mixed by pipetting up and down. An equal volume of 2X HBS (274 mM NaCl, 10 mM KCl, 42 mM HEPES, 1.4 mM Na₂PO₄, 12 mM dextrose, pH 7.05, JRH grade) was added with a 2-mL pipette, and the solution was pipetted up and down three times. The DNA mixture was immediately added to the 293 cells, one drop at a time, evenly throughout the dish. The cells were then incubated in a humidified environment at 37° C in 5% CO₂ for six hours. A granular precipitate was visible in the transfected cell cultures. After six hours, the DNA mixture was removed from the cells, which were then provided with fresh cell culture medium and incubated for an additional 72 hours.

After 72 hours, the cells were lysed and then treated with nuclease to reduce residual cellular and plasmid DNA. After precipitation, rAAV virions were purified by two cycles of isopycnic centrifugation; fractions containing rAAV virions were pooled, dialysed, and concentrated. The concentrated virions were formulated, sterile filtered (0.22 µM) and aseptically filled into glass vials. Vector genomes were quantified by the “Real Time Quantitative Polymerase Chain Reaction” method (Real Time Quantitative PCR. Heid C.A., Stevens J., Livak K.J., and Williams P.M. 1996. Genome Research 6:986-994. Cold Spring Harbor Laboratory Press).

Recombinant AAV1-hF.IX virions were produced in an analogous manner to rAAV6-hF.IX virions, with a pRep1Cap1 AAV helper function vector used in place of the pRepCap6 AAV helper function vector.

5 Recombinant AAV-2 virions were produced with the pHLP19 helper function vector (described in U.S. Patent No. 6,001,650, supra), the pLadeno5 plasmid, and the rAAV2-hF.IX expression plasmid.

EXAMPLE 2

HEMOPHILIA B TREATMENT IN RAG-1 MICE WITH 10 rAAV1-HF.IX, rAAV2-HF.IX, AND rAAV-6-HF.IX

15 RAG-1 female immunodeficient mice (homozygous for a mutation in the recombinase activating gene 1, and functionally equivalent to severe combined immunodeficiency mice because these mice do not produce mature B or T cells) 4-6 weeks old (obtained from Jackson Laboratories, Bar Harbor, ME) were injected with rAAV-6 virions (prepared as described in Example 1). Mice were anesthetized with an intraperitoneal injection of ketamine (70 mg/kg) and xylazine (10 mg/kg), and a 1 cm longitudinal incision was made in the lower extremity. Recombinant AAV6-hF.IX (2 x 10¹¹ viral vector genomes/kg in HEPES-Buffered-Saline, pH 7.8) virions were injected
20 into the tibialis anterior (25 µL) and the quadriceps muscle (50 µL) of each leg using a Hamilton syringe. Incisions were closed with 4-0 Vicryl suture. Blood samples were collected at seven-day intervals from the retro-orbital plexus in microhematocrit capillary tubes and plasma assayed for hF.IX by ELISA. Human F.IX antigen in mouse plasma was assessed by ELISA as described by Walter et al. (*Proc Natl Acad Sci USA* 25 (1996) 3:3056-3061). The ELISA did not cross-react with mouse F.IX. All samples were assessed in duplicate. Protein extracts obtained from injected mouse muscle were prepared by maceration of muscle in PBS containing leupeptin (0.5 mg/mL) followed by sonication. Cell debris was removed by microcentrifugation, and 1:10 dilutions of the protein extracts were assayed for hF.IX in the ELISA. The circulating plasma
30 concentrations of hF.IX, as measured by ELISA after three weeks post-IM injection, were 185 ng/mL for rAAV-6 hF.IX gene delivery, 110 ng/mL for rAAV1-hF.IX gene

delivery, and 6 ng/mL for rAAV-2 hF.IX gene delivery. After seven weeks post-injection, hF.IX plasma concentrations increased to approximately 190 ng/mL for rAAV6- hF.IX gene delivery, 200 ng/mL for rAAV1-hF.IX gene delivery, and 20 ng/mL for rAAV2-hF.IX gene delivery (see FIG. 1). After eleven weeks post-injection, hF.IX plasma concentrations increased to approximately 300 ng/mL for rAAV1-hF.IX gene delivery, but decreased to approximately 10 ng/mL for rAAV2-hF.IX gene delivery.

EXAMPLE 3

HEMOPHILIA B TREATMENT IN DOGS WITH AAV1-cF.IX

A colony of dogs having severe *hemophilia* B comprising males that are hemizygous and females that are homozygous for a point mutation in the catalytic domain of the canine factor IX (cF.IX) gene, was used to test the efficacy of cF.IX delivered by rAAV-1 virions (rAAV1-cF.IX). The severe hemophilic dogs lack plasma cF.IX, which results in an increase in whole blood clotting time (WBCT) to >60 minutes (normal dogs have a WBCT between 6-8 minutes), and an increase in activated partial thromboplastin time (aPTT) to 50-80 seconds (normal dogs have an aPTT between 13-18 seconds). These dogs experience recurrent spontaneous hemorrhages. Typically, significant bleeding episodes are successfully managed by the single intravenous infusion of 10 mL/kg of normal canine plasma; occasionally, repeat infusions are required to control bleeding.

Under general anesthesia, hemophilia B dogs were injected intramuscularly with rAAV1-cF.IX virions at a dose of 1×10^{12} vg/kg. The animals were not given normal canine plasma during the procedure.

Whole blood clotting time was assessed for cF.IX in plasma. Activated partial thromboplastin time was measured. A coagulation inhibitor screen was also performed. Plasma obtained from a treated hemophilic dog and from a normal dog was mixed in equal volumes and was incubated for 2 hours at 37° C. The inhibitor screen was scored as positive if the aPTT clotting time was 3 seconds longer than that of the controls (normal dog plasma incubated with imidazole buffer and pre-treatment hemophilic dog

plasma incubated with normal dog plasma). Neutralizing antibody titer against AAV vector was assessed.

In the hemophilia B dogs injected with AAV1-cF.IX, WBCT was shortened from > 60 min to 13 min (normal: 12-15 min).

5

EXAMPLE 4

HEMOPHILIA B TREATMENT IN HUMANS WITH AAV6-HF.IX

10 On Day 0 of the protocol patients are infused with hF.IX concentrate to bring factor levels up to ~100%, and, under ultrasound guidance, rAAV6-h.FIX virions are injected directly into 10-12 sites in the *vastus lateralis* of either or both anterior thighs. Injectate volume at each site is 250-500 μ L, and sites are at least 2 cm apart. Local anesthesia to the skin is provided by ethyl chloride or eutectic mixture of local
15 anesthetics. To facilitate subsequent muscle biopsy, the skin overlying several injection sites is tattooed and the injection coordinates recorded by ultrasound. Patients are observed in the hospital for 24 h after injection; routine isolation precautions will be observed during this period to minimize any risk of horizontal transmission of virions. Patients are discharged and seen daily in outpatient clinic daily for three days after
20 discharge, then weekly at the home hemophilia center for the next eight weeks, then twice monthly up to five months, then monthly for the remainder of the year, then annually in follow-up. Circulating plasma levels of hF.IX are quantified using ELISA as described in Example 2.